

Research  
Tools

Reviews

Journal  
CollectionNews &  
CommentBooks &  
Labware

Help



Feed

journals.bmn.com

Latest  
Updates

A-Z list MEDLINE Search



My BMN



Exit

Quick Site Search



Advanced site search

Jump to :

Vol

Page



Fill in Vol and Page

QUESTIONNAIRE

Questionnaire

Give us your opinion...

## Current Opinion in Chemical Biology

[Full A-Z Journal List](#) > Abstract

### Solid support linker strategies

[Review article]

Bradley J Backes, Jonathan A Ellman

*Current Opinion in Chemical Biology* 1997, 1:86-93.

#### Abstract:

The selection of an appropriate linker is critical to the success of any strategy for the solid-phase synthesis of small molecule libraries. While the primary function of the linker is to covalently attach the initial substrate to the support, innovative strategies have emerged recently in which linkers fulfill important auxiliary roles. These include the cleavage of compounds into solution leaving no trace of the support attachment site, cleavage via cyclization, cleavage by introduction of additional diversity into the structure, and cleavage whereby portions of the compound are sequentially released into solution.

Full text: [text](#), [+ thumbnails](#), [+ full figures](#), PDF 149 k

BioMedNet

Research  
Tools

Reviews

Journal  
CollectionNews &  
CommentBooks &  
LabwareScience  
JobsWeb  
Links
[Information for Advertisers](#)

© Elsevier Science Limited 2000

# Solid support linker strategies

Bradley J Backes\* and Jonathan A Ellman†

The selection of an appropriate linker is critical to the success of any strategy for the solid-phase synthesis of small molecule libraries. While the primary function of the linker is to covalently attach the initial substrate to the support, innovative strategies have emerged recently in which linkers fulfill important auxiliary roles. These include the cleavage of compounds into solution leaving no trace of the support attachment site, cleavage via cyclization, cleavage by introduction of additional diversity into the structure, and cleavage whereby portions of the compound are sequentially released into solution.

## Addresses

Department of Chemistry, University of California at Berkeley, Berkeley CA 94702, USA

\*e-mail: backes@uclink4.berkeley.edu

†e-mail: jellman@uclink.berkeley.edu

Current Opinion in Chemical Biology 1997, 1:86–93

<http://biomednet.com/elecreff/1367593100100086>

© Current Biology Ltd ISSN 1367-5931

## Abbreviations

DBU	1,8-diazabicyclo[5.4.0]undec-7-ene
DMF	<i>N,N</i> -dimethyl formamide
HF	hydrofluoric acid
TFA	trifluoroacetic acid
THF	tetrahydrofuran
THP	tetrahydropyranyl ether

## Introduction

Combinatorial technologies encompass numerous strategies to prepare and screen collections or 'libraries' of structurally related molecules [1,2]. Although library approaches have had an impact on the fields of materials development, catalysis, and molecular recognition [3], the major application has been the development of new therapeutic agents. The focus of these efforts has shifted to small molecule libraries [4\*,5\*] from peptide and oligonucleotide libraries since these biopolymers typically suffer from poor oral availability and rapid *in vivo* clearance.

General and high yielding synthesis methods must be devised in order to prepare structurally and functionally diverse small molecule libraries. While several solution-phase methods have been developed [4\*,5\*], to date the majority of library synthesis efforts have employed the solid-phase strategies pioneered by Merrifield for peptide synthesis [6]. The methods for substrate attachment to the support and the final cleavage of the products into solution are crucial features of all solid-phase strategies. The primary function of the linker is simply to covalently attach the substrate to the support. However, innovative

linkage strategies have emerged that allow for the cleavage of compounds into solution leaving no trace of the support attachment site, cleavage via cyclization, and cleavage by introduction of additional diversity into the structure. Linkers can also fulfill auxiliary roles in the synthesis of optically pure compounds and in the deconvolution of assay mixtures.

The following review will provide an overview of new linkage strategies for small molecule synthesis and combinatorial library applications from 1992 to the present, focusing primarily on the advances of the past two years. Many carboxylic acid- and amide-based linkers utilized for solid-phase peptide synthesis [7] have been adapted to small molecule synthesis. These developments have been reviewed elsewhere [4\*,5\*] and will not be discussed.

## General considerations for library synthesis

In solid-phase synthesis strategies, a substrate is attached to a functionalized polymer. Reactive polymers are prepared using copolymerization processes that incorporate functionalized monomers, or by direct derivatization of the polymer itself [8]. A functionalized polymer is typically coupled to a linker compound or 'handle' that provides the desired site for attachment of the substrate [9]. 'Preformed handle' strategies are those in which the handle is attached to the substrate in solution, and subsequently coupled to the polymer. Preformed handle strategies are effective when attaching a template or scaffold upon which diversity is introduced. Direct loading of the substrate onto the resin-bound linker is necessary when attaching multiple substrate building blocks.


For the solid-phase synthesis of small molecules, the choice of a linkage strategy deserves special consideration because linker stability and cleavage conditions dictate the range of synthetic chemistries that can be employed. A successful linkage strategy has the same requirements as a solution-phase protecting group strategy: quantitative loading (protection), chemical stability during the synthesis sequence (orthogonality), and quantitative cleavage (deprotection) without product degradation. Synthesis in a library format has the additional requirement that the cleavage conditions employed should not complicate rapid compound isolation.

## Protecting-group linker strategies

For most library synthesis efforts the initial building block or scaffold bears functionality (carboxylic acid, phenol, alcohol, amine etc.) that is covalently attached to a support-bound protecting group. After the structure is elaborated, the target compound is cleaved, bearing the same functionality. The development of new protecting-group linkers is certain to provide researchers with more

flexibility when planning a synthesis strategy. Only a brief summary of new advances will be given since these strategies are conceptually similar to their solution-phase counterparts.

Figure 1



Protecting-group linkers. Linkers used to attach compounds to the support through functionality present in the initial substrate and the final library products. A preformed handle approach has been employed with linkers marked with an \*.

Figure 1 shows linkers used in protecting-group linker strategies. All bold numbers refer to the structures shown in the figures. In early solid-phase synthesis reports, ester [10] and trityl ether [11,12] functionalities served to tether alcohols to a support. More recently, tetrahydropyranyl ether- (THP) 1 [13], acetal- 2 [14,15], ketal- 3 [16], silyl ether- 4, 5 [16,17] as well as carbonate-based linkages [18] have been utilized. For phenol attachment, Wang resin 6 is routinely employed to provide a *p*-alkoxybenzyl ether linkage [19]. Acid-labile benzyl carbamate based linkers for tethering amines to support were originally developed for amino→carboxyl peptide synthesis [20,21]. Functionalized resins 8 [22], 9 [23], and 10 [24] have recently been employed to provide a more labile *p*-alkoxybenzyl carbamate linkage. A palladium-labile allyl carbamate-based linkage has also been described [25]. The

ADCC 1° amine protecting group inspired the preparation of resin 11 [26]; the linkage is stable to trifluoroacetic acid (TFA) and stable to the bases piperidine and DBU, but is readily cleaved by 2% hydrazine in DMF. The THP linker 1 described above has also been used for attaching 2,6-dichloropurine to a support through the N-9 nitrogen [27]. Resin 12 [28] and the commercially available Seiber resin 13 [29] have been reductively aminated employing amines and aldehydes, respectively, to give support-bound amines. Additionally, trityl chloride resins are commonly employed for anchoring amines, diamines and amino alcohols to supports [30].


Linkage through pharmacophore-based functional groups is particularly useful because this site of a molecule is not varied in a library synthesis [31]. Alkoxyamine resins 14 [32,33] and 15 [32] allow for the synthesis of the hydroxamic acid functionality, a pharmacophore for metalloprotease inhibition. Rink 16 [34], Wang 6 [35], and Sasrin 7 [35] resins have all been utilized to provide sulfonamide-based linkages, and a semicarbazone-based linker (17) has been developed for anchoring amino acid aldehydes to support [36]. In addition, a phosphonate ester-based linkage has been employed for the synthesis of  $\alpha$ -amino [37] and  $\alpha$ -hydroxy phosphonates [38].

### Traceless linker strategies

Protecting-group linkage strategies are most useful when the unmasked functionality is important for the biological activity of the target compounds. Since auxiliary polar functionality can adversely affect the pharmacokinetic properties or binding properties of the target compounds, 'traceless' linker strategies have been devised that leave the target compound with no 'memory' of solid-phase synthesis (Fig. 2). Several metal-based aryl linkers have been described that permit cleavage with electrophilic reagents, often introducing additional diversity. In an early report, benzodiazepines were prepared using an aminoaryl stannane attached to the support through a silyl linkage 18 [39]. Treatment with HF provides benzodiazepines (shown as 22) free of auxiliary functionality. A more labile germanium derivative (19) has been prepared for the same purpose [40]. Arylsilane-based linkers 24 [41] and 25 [42\*] and a silyl ether based linker 26 [43] have been reported as well. Electrophilic cleavage employing Br<sub>2</sub> [40,42\*] and ICl [42\*] to provide bromine and iodine substituted aromatic products has been demonstrated in addition to protodesilylation by acidolysis [40,41,42\*] or basic fluoridolysis [41,43]. All of the reported metal-based aryl linkers utilize preformed handle strategies. It is likely that new strategies will be developed that provide more efficient loading of the substrate onto the solid support.

Traceless strategies that do not incorporate metals include thioether- 27 [44], 28 [45], 29 [45] and triazine- 30 [46] based linkers. For all of the strategies described in this section, the linkage group is replaced by either a hydrogen or halogen substituent, thus leaving a subtle

Figure 2



Traceless linkers. Linkers used to cleave compounds into solution leaving no trace of the support attachment site.


mark of attachment. A true traceless linkage strategy has been developed for the synthesis of isoquinolines [47<sup>\*</sup>]. Hydrolysis of Reissert complex 34 results in elimination of cyanide to provide isoquinoline products devoid of any evidence of solid-phase synthesis.

### Release by cyclization strategies

The release of compounds into solution by cyclization, whereby a reactive functionality closes down upon the support attachment site, provides a useful cleavage method. Linkers used in these strategies are shown in Figure 3. Typically, an unmasked support-bound nucleophile closes down upon a carbamate- or ester-based linkage, releasing the target compound into solution. Naturally, these strategies are limited to compounds that possess rings. It should be noted that due to competing intersite reactions [48], release by cyclization of rings larger than 5–6 members can result in substrate dimerization and resin cross-linking [49]. While these strategies can provide products in high purity when support-bound deletion sequences do not cleave into solution, a high fidelity synthetic sequence is still required to obtain uniform yields. Amino acids linked through a C-terminal ester linkage (36) have been employed for the synthesis of benzodiazepines 38 and hydantoins 40

40 with acid-promoted cleavage [50]. Hydantoins have also been prepared using a base-promoted cyclization method employing a carbamate linkage 41 [51]. A recently reported strategy to provide 5-alkoxyhydantoins (43) permits the introduction of additional elements of diversity to the library after release into solution during the cleavage step [52]. Diketopiperazines (45) have been prepared [53,54] using a strategy that results in *in situ* release when R<sup>1</sup> substituents are not sterically demanding [53]. In addition, base-promoted cyclization strategies have been employed to prepare 3-substituted quinazoline-2,4-diones [55], 5,6-dihydropyrimidine-2,4-diones [56], and 1,4-benzodiazepine-2,5-diones [57]. Cyclizations have also been performed in solution after cleavage of the compounds from the support. High fidelity syntheses are especially important for solution cyclization strategies since deletion sequences or side products are also released into solution. Diketopiperazines [58], diketomorpholines [53], and  $\beta$ -turn mimetics [59] have been prepared in this manner.

Figure 3




Linkers used in release by cyclization. Synthesis strategies in which target compounds are released via cyclization.

### Diversification linker strategies

Linkage strategies in which diverse substituents are incorporated into the target compound upon cleavage

are particularly useful for combinatorial synthesis efforts. Figure 4 details examples of these linkers. When synthesizing a library, it is desirable to employ limiting quantities of the cleavage agent, which results in products in solution that do not require further purification. The use of polymer-supported covalent scavengers [60\*] and sequestering agents [59,61] may provide access to pure products even when excess cleavage reagents are used. In an early report, a method was described to transesterify benzyl ester-based linkages with various alcohols under relatively mild conditions (LiBr and DBU in THF)[62]. Benzyl ester-based linkages have also been cleaved by aminolysis to give a range of amide products using aluminum trichloride [63] and trimethyl aluminum [64] reagents to promote the reaction. The harsh conditions employed and reagent byproducts produced may limit the generality of these methods. More reactive esters prepared from commercially available Kaiser's oxime resin [65,66] can be cleaved under mild conditions with amine nucleophiles to provide amide products in solution [67].

Figure 4



Diversification linkers. Linkers used to incorporate diverse substituents into the target compounds in the cleavage steps.

High reactivity towards cleavage limits the range of chemistry that can be employed during library synthesis. To address this problem, 'safety-catch' linkers that provide a more chemically stable linkage have been developed. A chemoselective activation step after solid-phase synthesis is complete provides a reactive linkage element for cleavage. Kenner's arylsulfonamide safety-catch linker, 46, provides a carboxylic acid linkage stable to both acid and base [68,69]. Activation via *N*-alkylation with


diazomethane and cleavage with hydrazine, amine and alkoxy nucleophiles gives hydrazide, amine, ester, and acid products in solution. Recently, a haloacetonitrile activation method has been reported that provides a highly reactive linkage for cleavage [70\*]. Limiting amounts of nucleophilic amines can be added to the support-bound *N*-cyanomethyl acylsulfonamide, resulting in complete conversion to provide pure amide products in solution. In order to apply this activation method to benzoic acid and carboxylic acids that possess  $\alpha$ -electron withdrawing groups such as amino acids, a second alkyl sulfonamide linker (47) was prepared. A sulfur-based safety-catch linker (48) prepared by direct loading of a thiol resin has been employed for the synthesis of small heterocycles [71\*]. After an oxidative activation step, nucleophilic cleavage with limiting amounts of various amines provides substituted heterocycles. A safety-catch linker for the synthesis of 3° amines has also been described [72\*]. Michael addition of an amine to acrylate ester resin (REM linker) 49 provides a support-bound 3° amine (50) for further elaboration. Quaternization with a variety of reactive alkylating reagents introduces diversity, and at the same time provides a labile linkage. Hoffman elimination with Hunig's base gives relatively pure 3° amines and regenerates the REM linker.

Additional diversification linkage strategies that do not involve an activation step have also been reported. Phthalamide resin 53 has been employed for the synthesis of phthalhydrazides [73]. Cleavage of 53 with substituted hydrazides provide the corresponding phthalhydrazide products. In preliminary studies, the addition of Grignard reagents to Weinreb amide linker 54 was employed to provide ketone products [74]. A related Weinreb amide linker was previously reported for the synthesis of  $\alpha$ -*N*-acylamino aldehydes [75]. Treatment of alkyl sulfonyl chloride resin 55 [76] with alcohols provides mesylate-based linkages to the support. Mesylate displacement with NaI has been demonstrated to afford alkyl iodide products. The authors of [75] suggest that the mesyl linkage could be used for a wide range of substitution reactions.

### Support-bound chiral auxiliaries

Polymer-supported chiral auxiliaries [77] for carrying out diastereoselective transformations to provide enantiomer-enriched compounds upon cleavage have been reported (representative examples are shown in Fig. 5). Since chirality is an important determinant in receptor recognition, this class of linkers could be particularly useful for the preparation of lead optimization libraries. Kurth and co-workers utilized polymer-bound auxiliary 56 [78] and  $C_2$ -symmetric auxiliary 57 [79] for preparing optically active 3,5-disubstituted- $\gamma$ -butyrolactones. In addition, a reusable polymer-bound 'Evans' oxazolidinone (58) has been reported [80]. Diastereoselective enolate alkylation was demonstrated with hydrolytic cleavage to provide enantiomerically-enriched  $\alpha$ -alkylated carboxylic acids [80].

Figure 5



Support-bound chiral auxiliaries.


### Photolabile linkers

Photolabile linkers [7] deserve special consideration since photolysis can provide an orthogonal and noninvasive cleavage strategy allowing for the direct assay of the cleaved library. Figure 6 shows examples of photolabile linkers. Photolabile linkers based upon the nitrobenzyl group, represented by 59, have been used extensively to provide ester- and amide-based linkages to the support. Related photolabile anchors have been employed for tethering alcohols [18,81–83] and amines [84] to support, mainly through carbonate and carbamate linkages, respectively. Photolysis liberates the target compound, resulting in the formation of a support-bound nitroso aldehyde. This functionality can trap liberated compounds, for example by reaction with 1° amines to give support-bound imines. Accordingly, photolabile linkers 60 [85], 61 [86], 62 [87], which provide a less reactive nitroso ketone functionality upon photolysis, have been developed. The nitroso byproducts can dimerize to form azo compounds that act as an internal light filter which retards the rate of cleavage. Methoxy substituents have been introduced to nitrobenzyl linkers to provide nitroveratryl-based linkers (63), since nitroveratryl groups have much higher absorptivity at the excitation wavelength than do nitrobenzyl groups. It should be noted, however, that Krafft *et al.* [88] have determined that the quantum yield for photodeprotection of nitroveratryl groups can be much lower than for nitrobenzyl groups, resulting in comparable cleavage photoefficiencies. One major limitation of photolabile linkers is chemical instability. The nitro functionality is unstable to most reducing agents and many organometallic reagents. In fact, the issue of chemical instability has inspired the recent development of a safety-catch photolabile linker (64) [89] that is activated towards photolytic cleavage by thioketal deprotection.

### Linkers for deconvolution

For some deconvolution approaches, that is to say the evaluation of progressively smaller pools of compounds to identify the active compound, partial release of the support-bound compound into solution is required. Photolabile linkers are ideal for the controlled partial release of compounds into solution (Fig. 7). As demonstrated for photolabile linker 65, long-wavelength UV light facilitates partial release, allowing multiple determinations to be


Figure 6



Photolabile linkers. Linkers that permit photolytic cleavage.

made from a single bead in conjunction with solution assays [82].

Figure 7



Linkers for deconvolution. Linkers used for the partial cleavage of compounds into solution for deconvolution strategies.

A second strategy for partial compound release is to employ multiple linkers that are cleaved under different reaction conditions. Multiple release linker 66 [90], which has been used for deconvolution studies to identify active peptides, relies on orthogonal cleavage conditions. One third of the peptide strands are cleaved at neutral pH via diketopiperazine formation, and one third are released at high pH via saponification. The identity of

the remaining support-bound peptide strand is determined by microsequencing. A related doubly-cleavable linker for peptide synthesis (67) has also been reported [91]. A portion of the support-bound compound is cleaved via diketopiperazine formation after protecting group removal with TFA. A second portion can then be cleaved under basic conditions. Multiple linkers have also been used in which compound release is governed by the relative acid lability of the linkers 68 [92]. It should be noted that multiple linker strategies drastically limit the range of chemistry that can be performed in a synthesis sequence since all synthesis steps must be orthogonal to the different cleavage strategies employed. The general utility of multiple release linkers for the synthesis of small molecule libraries is therefore likely to be limited.

## Conclusions

The methods for compound attachment to the support and cleavage into solution are crucial features of all solid-phase strategies. While the primary function of a linker is to covalently attach the initial substrate to a support, general approaches have been developed in which linkers fulfill important auxiliary roles. In this review we have described general strategies for four cleavage types: the cleavage of compounds into solution leaving no trace of the support attachment site; cleavage via cyclization; cleavage by introduction of additional diversity into the structure; and cleavage whereby portions of the compound are sequentially released into solution. An increasing number of researchers are applying library strategies to a host of research problems in chemistry and biology. The development of new linkage strategies will be driven by the need to prepare libraries of new compound classes as required to address specific research problems. A majority of these new linkage strategies will fall within one of the designated linker categories.

## Acknowledgements

JAE acknowledges the support of the National Science Foundation and the National Institutes of Health. BJB gratefully acknowledges the American Chemical Society Division of Organic Chemistry and Abbott Laboratories for predoctoral fellowship support.

## References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
  - of outstanding interest
- Gallop MA, Barrett RW, Dower WJ, Fodor SPA, Gordon EM: Applications of combinatorial technologies to drug discovery. 1. Background and peptide combinatorial libraries. *J Med Chem* 1994, 37:1233-1251.
  - Gordon EM, Barrett RW, Dower WJ, Fodor SPA, Gallop MA: Applications of combinatorial technologies to drug discovery. 2. Combinatorial organic synthesis, library screening strategies, and future directions. *J Med Chem* 1994, 37:1385-1401.
  - Czarnik AW, Ellman JA (Eds.): Special issue: combinatorial chemistry. *Accounts Chem Res* 1996, 29:112-170. This issue includes articles describing several applications of library strategies, including materials science, molecular recognition, and therapeutics.
  - Thompson LA, Ellman JA: Synthesis and applications of small molecule libraries. *Chem Rev* 1996, 96:555-600.
  - Library synthesis and evaluation strategies, as well as solution- and solid-phase methods to prepare small compound libraries, are reviewed.
  - Balkenhohl F, von dem Bussche-Hünnefeld C, Lansky A, Zechel C: Combinatorial synthesis of small molecule libraries. *Angew Chem Int Ed* 1996, 35:2288-2337.
  - Library synthesis and evaluation strategies, as well as solution- and solid-phase methods to prepare small compound libraries, are reviewed.
  - Merrifield RB: Solid phase peptide synthesis I. The synthesis of a tetrapeptide. *J Am Chem Soc* 1963, 85:2149-2150.
  - Lloyd-Williams P, Albericio F, Giralt E: Convergent solid-phase peptide synthesis. *Tetrahedron* 1993, 49:11065-11133.
  - Barany G, Merrifield RB: Solid phase peptide synthesis. In *The Peptides*. Edited by Gross E, Meienhofer J. New York: Academic Press, Inc; 1980:43-55.
  - Keifer PA: Influence of resin structure, tether length, and solvent upon the high-resolution  $^1\text{H}$  NMR spectra of solid-phase synthesis resins. *J Org Chem* 1996, 61:1558-1559.
  - The quality of  $^1\text{H}$  NMR spectra of support-bound compounds using different combinations of resins, linkers and solvents were evaluated. The findings suggest that polymer and linker combinations that provide a longer tether, and therefore increased substrate mobility, provide superior spectra.
  - Leznoff CC, Wong JY: The use of polymer supports in organic synthesis. The synthesis of monotrityl ethers of symmetrical diols. *Can J Chem* 1972, 50:2892-2893.
  - Fréchet JM, Nuyens LJ: The use of polymers as protecting groups in organic synthesis. III. Selective functionalization of polyhydroxyl alcohols. *Can J Chem* 1976, 54:926-934.
  - Fyles TM, Leznoff CC: The use of polymer supports in organic synthesis. V. The preparation of monoacetates of symmetrical diols. *Can J Chem* 1976, 54:935-942.
  - Thompson LA, Ellman JA: Straightforward and general method for coupling alcohols to solid-supports. *Tetrahedron Lett* 1994, 35:9333-9336.
  - Wang GT, Li S, Wideburg N, Kraft GA, Kempf DJ: Synthetic chemical diversity: solid phase synthesis of libraries of  $\text{C}_2$  symmetric inhibitors of HIV protease containing diamino diol and diamino alcohol cores. *J Med Chem* 1995, 38:2995-3002.
  - Wong JY, Manning C, Leznoff CC: Solid phase synthesis and photochemistry of 4,4'-stilbenedicarbaldehyde. *Angew Chem Int Ed* 1974, 13:666-667.
  - Randolph JT, McClure KF, Danishefsky SJ: Major simplifications in oligosaccharide syntheses arising from a solid-phase based method: an application to the synthesis of the Lewis b antigen. *J Am Chem Soc* 1995, 117:5712-5719.
  - Chan TH, Huang WQ: Polymer-anchored organosilyl protecting group in organic synthesis. *J Chem Soc Chem Commun* 1985:909-911.
  - Alsina J, Chiva C, Ortiz M, Rabanal F, Giralt E, Albericio F: Active carbonate resins for solid-phase synthesis through the anchoring of a hydroxyl function. Synthesis of cyclic and alcohol peptides. *Tetrahedron Lett* 1997, 38:883-886.
  - Sarshar S, Siev D, Mjalli AMM: Imidazole libraries on solid support. *Tetrahedron Lett* 1996, 37:835-838.
  - Letsinger RL, Kornet MJ: Popcorn polymer as a support in multistep syntheses. *J Am Chem Soc* 1963, 85:3045-3046.
  - Felix AM, Merrifield RB: Azide solid phase peptide synthesis. *J Am Chem Soc* 1970, 92:1385-1391.
  - Hauske JR, Dorff P: A solid phase CBZ chloride equivalent - a new matrix specific linker. *Tetrahedron Lett* 1995, 36:1589-1592.
  - Kaljuste K, Undén A: Solid phase synthesis of 1,2,3,4-tetrahydro- $\beta$ -carboline; implications for combinatorial chemistry. *Tetrahedron Lett* 1995, 36:9211-9214.
  - Alsina J, Rabanal F, Giralt E, Albericio F: Solid-phase synthesis of "head-to-tail" cyclic peptides via lysine side-chain anchoring. *Tetrahedron Lett* 1994, 35:9633-9636.
  - Kaljuste K, Undén A: Solid-phase synthesis of peptide aminoalkylamides using an allyl linker. *Tetrahedron Lett* 1996, 37:3031-3034.

26. Bannwarth W, Huebscher J, Barner R: A new linker for primary amines applicable to combinatorial approaches. *Bioorg Med Chem Lett* 1996, 6:1525-1528.
27. Nugiel DA, Cornelius LAM, Corbett JW: Facile preparation of 2,6-disubstituted purines using solid-phase chemistry. *J Org Chem* 1997, 62:201-203.
28. Boojamara CG, Burrow KM, Ellman JA: An expedient and high-yielding method for the solid-phase synthesis of diverse 1,4-benzodiazepine-2,5-diones. *J Org Chem* 1995, 60:5742-5743.
29. Chan WC, Mellor SL: Reductive alkylation of 9-amino-xanthen-3-yloxymethylpoly(styrene): a novel procedure for the synthesis of peptidyl *N*-alkyl amides by Fmoc/Bu<sup>t</sup> chemistry. *J Chem Soc Chem Commun* 1995:1475-1476.
30. Krchnák V, Weichsel AS, Cabel D, Flegelova Z, Lebl M: Structurally homogeneous and heterogeneous synthetic combinatorial libraries. *Mol Divers* 1995, 1:149-164.
31. Kick EK, Ellman JA: Expedient method for the solid-phase synthesis of aspartic acid protease inhibitors directed toward the generation of libraries. *J Med Chem* 1995, 38:1427-1430.
32. Floyd CD, Lewis CN, Patel SR, Whittaker M: A method for the synthesis of hydroxamic acids on solid phase. *Tetrahedron Lett* 1996, 37:8045-8048.
33. Richter LS, Desai MC: A TFA-cleavable linkage for solid-phase synthesis of hydroxamic acids. *Tetrahedron Lett* 1997, 38:321-322.
34. Beaver KA, Siegmund AC, Spear KL: Application of the sulfonamide functional group as an anchor for solid phase organic synthesis (SPOS). *Tetrahedron Lett* 1996, 37:1145-1148.
35. Ngu K, Patel DV: Preparation of acid-labile resins with halide linkers and their utility in solid phase organic synthesis. *Tetrahedron Lett* 1997, 38:973-976.
36. Murphy AM, Dagnino R Jr, Vallar PL, Trippe AJ, Sherman SL, Lumpkin RH, Tamura SY, Webb TR: Automated synthesis of peptide C-terminal aldehydes. *J Am Chem Soc* 1992, 114:3156-3157.
37. Zhang C, Mjalli AMM: A combinatorial method for the solid phase synthesis of  $\alpha$ -amino phosphonates and phosphonic acids. *Tetrahedron Lett* 1996, 37:5457-5460.
38. Cao X, Mjalli AMM: Combinatorial method for the synthesis of  $\alpha$ -hydroxy phosphonates on Wang resin. *Tetrahedron Lett* 1996, 37:6073-6076.
39. Plunkett MJ, Ellman JA: A silicon-based linker for traceless solid-phase synthesis. *J Org Chem* 1995, 60:6006-6007.  
• An early demonstration of traceless synthesis employing an arylsilane-based linkage for benzodiazepine synthesis is described.
40. Plunkett MJ, Ellman JA: Germanium and silicon linking strategies for traceless solid-phase synthesis. *J Org Chem* 1997, in press.
41. Chenere B, Finkelstein JA, Veber DF: Protodetachable arylsilane polymer linkages for use in solid phase organic synthesis. *J Am Chem Soc* 1995, 117:11999-12000.
42. Han Y, Walker SD, Young RN: Silicon directed *ipso*-substitution of polymer bound arylsilanes: preparation of biaryls via the Suzuki cross-coupling reaction. *Tetrahedron Lett* 1996, 37:2703-2706.  
•  
Cleavage of an arylsilane-based linkage with several electrophiles, including Br<sub>2</sub>, ICl and TFA, to prepare bromine, iodine and hydrogen substituted aryl products is demonstrated.
43. Boehm TL, Showalter HDH: Development of a novel silyl ether linker for solid-phase organic synthesis. *J Org Chem* 1996, 61:6498-6499.
44. Sucholeiki I: Solid-phase photochemical C-S bond cleavage of thioethers - a new approach to the solid-phase production of non-peptide molecules. *Tetrahedron Lett* 1994, 35:7307-7310.
45. Zhao XY, Jung KW, Janda KD: Soluble polymer synthesis: an improved traceless linker methodology for aliphatic C-H bond formation. *Tetrahedron Lett* 1997, 38:977-980.
46. Nelson JC, Young JK, Moore JS: Solid-phase synthesis of phenylacetylene oligomers utilizing a novel 3-propyl-3-(benzyl-supported) triazene linkage. *J Org Chem* 1996, 61:8160-8168.
47. Lorschach BA, Müller RB, Kurth MJ: Reissert-based 'traceless' solid-phase synthesis: isoquinoline, and isoxazoline-containing heterocycles. *J Org Chem* 1996, 61:8716-8717.  
•  
A novel strategy for traceless synthesis is reported. The Reissert cleavage provides isoquinoline products that bear no evidence of solid-phase synthesis.
48. Bhargava KK, Sarin VK, Trang NL, Cerami A, Merrifield RB: Synthesis of a cyclic analogue of oxidized glutathione by an intersite reaction in a swollen polymer network. *J Am Chem Soc* 1983, 105:3247-3251.
49. Van Maarseveen JHV, den Hartog JAJ, Engelen V, Finner E, Visser G, Kruse CG: Solid phase ring-closing metathesis: cyclization/cleavage approach towards a seven membered cycloolefin. *Tetrahedron Lett* 1996, 37:8249-8252.
50. DeWitt SH, Kiely JS, Stankovic CJ, Schroeder MC, Reynolds-Cody DM, Pavia MR: "Diversomers": an approach to nonpeptide, nonoligomeric chemical diversity. *Proc Natl Acad Sci USA* 1993, 90:6909-6913.
51. Dressman BA, Spangle LA, Kaldor SW: Solid phase synthesis of hydantoins using a carbamate linker and a novel cyclization/cleavage step. *Tetrahedron Lett* 1996, 37:937-940.
52. Hanessian S, Yang RY: Solution and solid phase synthesis of 5-alkoxyhydantoin libraries with a three-fold functional display. *Tetrahedron Lett* 1996, 37:5835-5838.
53. Scott BO, Siegmund AC, Marlowe CK, Pei Y, Spear KL: Solid phase organic synthesis (SPOS): a novel route to diketopiperazines and diketomorpholines. *Mol Divers* 1995, 1:125-134.
54. Goodfellow VS, Laudeman CP, Gerrity JJ, Burkard M, Strobel E, Zuzack JS, McLeod DA: Rationally designed non-peptides: variously substituted piperazine libraries for the discovery of bradykinin antagonists and other G-protein-coupled receptor ligands. *Mol Divers* 1996, 2:97-102.
55. Gouilleux L, Fehrentz JA, Wintemitz F, Martinez J: Solid phase synthesis of chiral 3-substituted quinazoline-2,4-diones. *Tetrahedron Lett* 1996, 37:7031-7034.
56. Kolodziej SA, Hamper BC: Solid-phase synthesis of 5,6-dihydropyrimidine-2,4-diones. *Tetrahedron Lett* 1996, 37:5257-5280.
57. Mayer JP, Zhang J, Bjergaard K, Lenz DM, Gaudino JJ: Solid phase synthesis of 1,4-benzodiazepine-2,5-diones. *Tetrahedron Lett* 1996, 37:8081-8084.
58. Gordon DW, Steele J: Reductive alkylation on a solid phase: synthesis of a piperazinedione combinatorial library. *Bioorg Med Chem Lett* 1995, 5:47-50.
59. Virgilio AA, Schürer SC, Ellman JA: Expedient solid-phase synthesis of putative  $\beta$ -turn mimetics incorporating the i + 1, i + 2, and i + 3 sidechains. *Tetrahedron Lett* 1996, 37:6961-6964.
60. Kaldor SW, Siegel MG, Fritz JE, Dressman BA, Hahn PJ: Use of solid supported nucleophiles and electrophiles for the purification of non-peptide small molecule libraries. *Tetrahedron Lett* 1996, 37:7193-7196.  
•  
Support-bound nucleophiles and electrophiles are used to 'quench' excess reactants employed for amine alkylations and acylations performed in solution. Solvent filtration and evaporation provides products in high yield.
61. Gayo LM, Suto MJ: Ion-exchange resins for solution phase parallel synthesis of chemical libraries. *Tetrahedron Lett* 1997, 38:513-516.
62. Seebach D, Thaler A, Blaser D, Ko SY: 104. Transesterifications with 1,8-diazabicyclo[5.4.0]undec-7-ene/lithium bromide (DBU/LiBr) - also applicable to cleavage of peptides from resins in Merrifield syntheses. *Helv Chim Acta* 1991, 74:1102-1118.
63. Bam DR, Morphy JR, Rees DC: Synthesis of an array of amides by aluminium chloride assisted cleavage of resin-bound esters. *Tetrahedron Lett* 1996, 37:3213-3216.
64. Ley SV, Mynett DM, Koot WJ: Solid phase synthesis of bicyclo [2.2.2]octane derivatives via tandem Michael addition reactions and subsequent reductive amination. *Synlett* 1995:1017-1020.
65. DeGrado WF, Kaiser ET: Polymer-bound oxime esters as supports for solid-phase peptide synthesis. Preparation of protected peptide fragments. *J Org Chem* 1980, 45:1295-1300.
66. Mohan R, Chou YL, Morrissey MM: Pictet-Spengler reaction on solid support: synthesis of 1,2,3,4-tetrahydro- $\beta$ -carboline libraries. *Tetrahedron Lett* 1996, 37:3963-3966.



67. Voyer N, Lavoie A, Pinette M, Bernier J: A convenient solid phase preparation of peptide substituted amides. *Tetrahedron Lett* 1994, 35:355–358.
68. Kenner GW, McDermott JR, Sheppard RC: The safety catch principle in solid phase synthesis. *J Chem Soc Chem Commun* 1971:636–637.
69. Backes BJ, Ellman JA: Carbon-carbon bond-forming methods on solid support. Utilization of Kenners 'safety-catch' linker. *J Am Chem Soc* 1994, 116:11171–11172.
70. Backes BJ, Virgilio AV, Ellman JA: Activation method to prepare a highly reactive acylsulfonamide 'safety-catch' linker for solid-phase synthesis. *J Am Chem Soc* 1996, 118:3055–3056.
- Reports an activation method that provides a highly reactive acylsulfonamide for cleavage. Treatment of the activated acylsulfonamide with limiting amounts of nucleophiles provides products in solution without further purification.
71. Gayo LM, Suto MJ: Traceless linker: oxidative activation and displacement of a sulfur-based linker. *Tetrahedron Lett* 1997, 38:211–214.
- A sulfur-based linker is described. After an oxidative activation step, limiting amounts of amine nucleophiles can be added to cleave the products from support. Small substituted heterocycles are provided in solution without the need for further purification.
72. Morphy JR, Rankovic Z, Rees DC: A novel linker strategy for solid-phase synthesis. *Tetrahedron Lett* 1996, 37:3209–3212.
- A reusable safety-catch linker is reported to prepare 3° amines in high purity.
73. Nielson J, Rasmussen PH: Implementation of a combinatorial cleavage and deprotection scheme. 1. Synthesis of phthalhydrazide libraries. *Tetrahedron Lett* 1996, 37:3351–3354.
74. Dihn TQ, Armstrong RW: Synthesis of ketones and aldehydes via reactions of Weinreb-type amides on solid support. *Tetrahedron Lett* 1996, 37:1161–1164.
75. Fehrentz JA, Paris M, Heitz A, Velek J, Liu CF, Winternitz F, Martínez J: Improved solid phase synthesis of C-terminal peptide aldehydes. *Tetrahedron Lett* 1995, 36:7871–7874.
76. Hunt JA, Roush WR: Solid-phase synthesis of 6-deoxyoligosaccharides. *J Am Chem Soc* 1996, 118:9998–9999.
77. Worster PM, McArthur CR, Leznoff CC: Asymmetric synthesis of 2-alkylcyclohexanones on solid phases. *Angew Chem Int Ed* 1979, 18:221–222.
78. Moon HS, Schore NE, Kurth MJ: A polymer-supported chiral auxiliary applied to the iodolactonization reaction: preparation of  $\gamma$ -butyrolactones. *J Am Chem Soc* 1992, 114:6088–6089.
79. Moon HS, Schore NE, Kurth MJ: A polymer-supported C<sub>2</sub>-symmetric chiral auxiliary: preparation of non-racemic 3,5-disubstituted- $\gamma$ -butyrolactones. *Tetrahedron Lett* 1994, 35:8915–8918.
80. Allin SM, Shuttleworth SJ: The preparation and first application of a polymer-supported 'Evans' oxazolidinone. *Tetrahedron Lett* 1996, 37:8023–8026.
81. Nicolau KC, Winsinger N, Pastor J, DeRoos F: A general and highly efficient solid phase synthesis of oligosaccharides. Total synthesis of a heptasaccharide phytoalexin elicitor (HPE). *J Am Chem Soc* 1997, 119:449–450.
82. Baldwin JJ, Burbaum JJ, Henderson I, Ohmeyer MHJ: Synthesis of a small molecule combinatorial library encoded with molecular tags. *J Am Chem Soc* 1995, 117:5589–5590.
83. Venkatesan H, Greenburg MM: Improved utility of photolabile solid phase synthesis supports for the synthesis of oligonucleotides containing the 3'-hydroxyl termini. *J Org Chem* 1996, 61:525–529.
84. Armstrong RW, Combs AP, Tempest PA, Brown SD, Keating TA: Multiple-component condensation strategies for combinatorial chemistry. *Accounts Chem Res* 1996, 29:123–131.
85. Holmes CP, Jones DG: Reagents for combinatorial organic synthesis: development of a new *o*-nitrobenzyl photolabile linker for solid phase synthesis. *J Org Chem* 1995, 60:2318–2319.
86. Ajayaghosh A, Pillai VNR: Solid-phase synthesis and C-terminal amidation of peptides using a photolabile *o*-nitrobenzylhydraminopolystyrene support. *Tetrahedron Lett* 1995, 36:777–780.
87. Brown BB, Wagner DS, Geysen HM: A single-bead decode strategy using electrospray ionization mass spectrometry and a new photolabile linker – 3-amino-3-(2-nitrophenyl)propionic acid. *Mol Divers* 1995, 1:4–12.
88. Krafft GA, Sutton WR, Cummings RT: Photoactive fluorophores. 3. Synthesis and photoactivation of fluorogenic difunctionalized fluoresceins. *J Am Chem Soc* 1988, 110:301–303.
89. Routledge A, Abell C, Balasubramanian S: The use of a dithiane protected benzoin photolabile safety catch linker for solid-phase synthesis. *Tetrahedron Lett* 1997, 38:1227–1230.
90. Salmon SE, Lam KS, Leble M, Kandola A, Khattri PS, Wade S, Pátek M, Kočíš P, Krchňák V, Thorpe D, Felder S: Discovery of biologically active peptides in random libraries: solution-phase testing after staged orthogonal release from resin beads. *Proc Natl Acad Sci USA* 1993, 90:11708–11712.
91. Kočíš P, Krchňák V, Lebl M: Symmetrical structure allowing the selective multiple release of a defined quantity of peptide from a single bead of polymeric support. *Tetrahedron Lett* 1993, 34:7251–7252.
92. Cardno M, Bradley M: A simple multiple release system for combinatorial library and peptide analysis. *Tetrahedron Lett* 1996, 37:135–138.